

One-pion-exchange final-state-interaction and the $p\bar{p}$ near threshold enhancement in $J/\psi \rightarrow \gamma p\bar{p}$ decays

B.S.Zou and H.C. Chiang

CCAST(World Lab.), P.O.Box 8730, Beijing 100080, China

Institute of High Energy Physics, The Chinese Academy of Sciences,
Beijing 100039, China*

Institute of Theoretical Physics, The Chinese Academy of Sciences,
Beijing 100080, China

and

Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator,
Lanzhou 730000, China

Abstract

For the $N\bar{N}$ system, the one-pion-exchange (OPE) interaction gives the largest attractive force for $N\bar{N}$ with isospin $I = 0$ and spin $S = 0$, while a near threshold enhancement was observed for $p\bar{p}$ with $I = 0$ and $S = 0$ in $J/\psi \rightarrow \gamma p\bar{p}$ decays. With a K-matrix approach, we find that the OPE final-state-interaction (FSI) makes an important contribution to the near-threshold enhancement in the $p\bar{p}$ mass spectrum in $J/\psi \rightarrow \gamma p\bar{p}$ decays.

PACS number(s): 14.40.Cs., 13.75.cs., 13.66.Bc.

*Mailing address

Recently the BES Collaboration has observed a near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum from the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$ [1]. The enhancement can be fitted with either an S- or P-wave Breit-Wigner resonance function. No similar structure is seen in $J/\psi \rightarrow \pi^0 p\bar{p}$ decays. In the S-wave case, the peak mass is below $2M_p$ around $M = 1859 MeV$ with a total width $\Gamma < 30 MeV$. These observations together with other similar results in the decays of B mesons [2] stimulate further investigations for the quasi-bound nuclear baryonium or multiquark resonance near the $2M_p$ threshold. What is the origin of the $p\bar{p}$ enhancement at $M_{p\bar{p}} \approx 2M_p$ in the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$? Datta et al. [3] describe the enhancement as the formation of a zero baryon number, “deuteron-like” singlet 1S_0 state. Does it come from any quasi-bound nuclear baryonium or multiquark resonance near the $2M_p$ threshold? In order to draw a conclusion we must study other dynamics which might affect the spectrum of the outgoing proton and antiproton.

The question of possible nucleon-antinucleon($N\bar{N}$) bound states was raised many years ago, in particular by Fermi and Yang [4]. In the sixties, explicit attempts were made to describe the spectrum of ordinary mesons as $N\bar{N}$ bound states. It was noticed [5], however, that the $N\bar{N}$ picture hardly reproduces the observed patterns of the meson spectrum. Encouraged by the evidence from many intriguing experimental investigations new types of mesons with a mass near the $N\bar{N}$ threshold and specific decay properties were proposed [6,7]. However, at the time where several candidate for baryonium were proposed, the quasi-nuclear approach was seriously challenged by a direct quark picture. Stimulated by the success of the quark models, exotic multiquark configurations were studied extensively [8]. The observation of the Pentaquark state [9] stimulates further searches for other multiquark bound states.

It was noticed [10–12] that for a multiquark or quasi-bound hadronic system close to its dissociation threshold, two hadrons will experience their long-range interaction, in particular the pion exchange. “Hadronic molecule” states might be formed. Due to its long range nature, the pion exchange plays a crucial role in achieving the binding of some configuration, especially for two hadrons in relative S-state. In a chiral unitary approach

it was also found [13] that to solve the coupled channel Bethe-Salpeter equations is crucial for explaining the observations in the meson-meson and meson-baryon interactions. Similarly, the outgoing proton and antiproton from the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$ will experience the long-range final-state interaction before they are detected. In order to better understand the nature of the experimental observation of the near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum from the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$ one has to evaluate the final-state-interaction (FSI) contribution to the invariant mass spectrum near $M_{p\bar{p}} \approx 2M_p$.

In this note, with the one-pion-exchange (OPE) potential between the proton and antiproton, we study the FSI of $p\bar{p}$ by the K-matrix approach for the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$.

It is well known [14] that for the NN system, the central OPE potential is attractive for $(S, I) = (0, 1)$ (deuteron) or $(S, I) = (1, 0)$, and repulsive for $S = I = 0$ (strong) or $S = I = 1$ (weak). For the $N\bar{N}$ system, the meson exchange interaction is related to the corresponding one for the NN system by the G-parity transformation, and the OPE potential gets an additional negative sign due to the negative G-parity of the pion. Hence the central OPE potential gives the largest attractive force for $N\bar{N}$ with $S = I = 0$. The attractive force is three times stronger than the corresponding one for the deuteron. The near threshold narrow enhancement observed in the $J/\psi \rightarrow \gamma p\bar{p}$ happens to have quantum numbers $S = I = 0$ preferred [1]. From the one-pion-exchange theory [14,15], the nucleon-antinucleon potential can be written as

$$V_{p\bar{p}}^\pi = \frac{C^{SI} f_\pi^2}{\vec{q}^2 + m_\pi^2} \quad (1)$$

with $C^{00} = -3$ for $S = I = 0$, $C^{11} = -1/3$ and $C^{10} = C^{01} = 1$. f_π is the πNN coupling constant $f_\pi^2/4\pi \simeq 0.08$ and m_π the mass of the π meson. \vec{q} is the 3-momentum transfer between the proton and antiproton. The $p\bar{p}$ with $I = S = L = 0$ from the radiative decays $J/\psi \rightarrow \gamma p\bar{p}$ will experience the largest attractive long-range OPE final-state-interaction. From the one-pion-exchange potential, in principle, one could calculate the two-body $N\bar{N}$ scattering amplitude by solving the Bethe-Salpeter equation

$$\bar{T} = V + VG\bar{T}. \quad (2)$$

Here G is the loop function of a proton and an antiproton propagators. It has been shown [16] that the K-matrix formulism provides an elegant way of expressing the unitarity of the S-matrix for the processes of the type $a + b \rightarrow c + d$. In the K-matrix approach the invariant S-wave $p\bar{p}$ scattering T-matrix can be expressed as

$$\bar{T} = \frac{K_s}{1 - iK_s\rho_{p\bar{p}}} \quad (3)$$

where $\rho_{p\bar{p}}$ is the phase space factor for the $p\bar{p}$ system

$$\rho_{p\bar{p}} = \frac{M_p^2 k}{\pi\sqrt{s}} \quad (4)$$

with s the invariant mass squared of the $p\bar{p}$ system and $k = \sqrt{s/4 - M_p^2}$ the momentum magnitude of the proton in the proton-antiproton c.m. system. Following an usual approach for the strong interaction in the K-matrix formalism [17,18], the K_s is taken as the S-wave projection of the $N\bar{N}$ potential, i.e.,

$$K_s = \frac{1}{4k^2} \int_{-4k^2}^0 dt V_{p\bar{p}}^\pi(t) \quad (5)$$

where $t = -\vec{q}^2$. For the $I = S = 0$ case, K_s can be easily evaluated from Eq.(5) as

$$K_s = -\frac{3f_\pi^2}{4k^2} \ln\left(1 + \frac{4k^2}{m_\pi^2}\right). \quad (6)$$

In this approach, by considering the OPE FSI of the proton and antiproton, the T-matrix for $J/\psi \rightarrow \gamma p\bar{p}(^1S_0)$ decays can be written as

$$T_{J/\psi \rightarrow \gamma p\bar{p}}^{(0)} = \frac{T_{J/\psi \rightarrow \gamma p\bar{p}}^{(0)}}{1 - i\rho_{p\bar{p}}K_s} = \frac{T_{J/\psi \rightarrow \gamma p\bar{p}}^{(0)}}{1 + i\frac{3M_p^2}{k\sqrt{s}}\frac{f_\pi^2}{4\pi}\ln\left(1 + \frac{4k^2}{m_\pi^2}\right)}. \quad (7)$$

Here $T_{J/\psi \rightarrow \gamma p\bar{p}}^{(0)}$ is T-matrix of the bare $J/\psi \rightarrow \gamma p\bar{p}(^1S_0)$ without considering the FSI. The conservation of parity and total angular momentum requires the orbital angular momentum between γ and the $p\bar{p}(^1S_0)$ to be $L = 1$, so that the $T^{(0)}$ is proportional to the momentum of the photon K_γ in the J/ψ rest system, i.e.,

$$T_{J/\psi \rightarrow \gamma p\bar{p}}^{(0)} = CK_\gamma. \quad (8)$$

In reality, C should be an s -dependent function. Here to illustrate the OPE FSI effect, we assume C as a constant. In Fig.(1) we show the T-matrix squared as a function of the invariant mass of the proton and antiproton for the $J/\psi \rightarrow \gamma p\bar{p}(^1S_0)$ process. The solid line corresponds to that with the FSI and the dashed line is that without the FSI. We find that the final-state-interaction has an important contribution to the $p\bar{p}$ enhancement near $M_{p\bar{p}} = 2M_p$ in $J/\psi \rightarrow \gamma p\bar{p}$ decays. Compared with plateau region well above threshold, the OPE FSI enhancement factor at the $p\bar{p}$ threshold is larger than 2. The phenomena of a narrow near-threshold peak due to the t-channel pion exchange is not new. For example, the striking narrow peak near $p\omega$ threshold in the $\gamma p \rightarrow \omega p$ process is found to be produced by the t-channel pion exchange [19].

It is well known that there is a very large production of two gluon system with $J^{PC} = 0^{-+}$ below $2M_p$ from the J/ψ radiative decays [20–26]. So C should at least have some broad resonance peaks below $2M_p$, which have not been well understood. It is quite possible that the interference of those components plus the narrow OPE FSI structure could explain the $p\bar{p}$ near threshold enhancement in the $J/\psi \rightarrow \gamma p\bar{p}$ process.

In $J/\psi \rightarrow \pi^0 p\bar{p}$ decays, however, because of the isospin conservation the isospin of the $p\bar{p}$ system must be 1. The $p\bar{p}$ interaction is either repulsive or one order of magnitude weaker than for the isoscalar $p\bar{p}(^1S_0)$ system. One should not find the near-threshold $p\bar{p}$ enhancement. In the decays of B mesons $B^0 \rightarrow D^0 p\bar{p}$ and $B^\pm \rightarrow K^\pm p\bar{p}$, the isospin of the $p\bar{p}$ system has isospin 0. The enhancement of the low-mass $p\bar{p}$ systems in B decays may also be understood by the FSI. The very narrow proton-antiproton atomic states observed by LEAR experiments [27] at $p\bar{p}$ threshold may also play some role in various narrow structure observed recently near $p\bar{p}$ threshold.

We would like to thank profs. Zhang Zong-Ye, Yu You-wen, Shen Peng-nian for stimulating discussions. This work is supported in part by the National Natural Science Foundation of China under grand Nos. 10225525, 10055003, 10075057, 10147208 and 19975053 and by the Chinese Academy of Sciences under project No. KJCX2-SW-N02.

FIGURES




FIG. 1. T-matrix squared with (solid line) and without (dashed line) OPE FSI, with an arbitrary normalization.

REFERENCES

- [1] J.Z.Bai et al., Phys. Rev. Lett. 91 , 022001 (2003).
- [2] K. Abe et al., Phys. Rev. Lett. 88, 181803 (2002);
K. Abe et al., Phys. Rev. Lett. 89, 1851802 (2002).
- [3] A. Datta et al., e-print arXiv:hep-ph/0306097.
- [4] E.Fermi and C.N. Yang, Phys. Rev. 76, 1739 (1949).
- [5] J.S. Ball et al., Phys. Rev. 142,1000(1966).
- [6] I.S. Shapiro, Phys. Rept. 35, 129(1978).
- [7] C.B. Dover and J.M. Richard, Ann. Phys. 121, 70(1979).
- [8] R.L. Jaffe, Phys. Rev. D17,1444(1978).
- [9] T. Nakano et al., Phys. Rev. Lett. 91,102002(2003);
V.V. Barmin et al., e-print arXiv: hep-ex/0304040;
S. Stepanyan et al.,e-print arXiv: hep-ex/0307018;
J.Barth wt al., e-print arXiv: hep-ex/0307083.
- [10] N.A.Tornqvist, Phys. Rev. Lett. 67, 556 (1991); Z. Phys. C61, 525 (1994); e-print
arXiv: hep-ph/0308277.
- [11] T.E.O.Ericson and G.Karl, Phys. Lett. B309, 426 (1993).
- [12] J.M. Richard, Nucl.Phys.Proc.Suppl. 86, 361(2000).
- [13] J.A. Oller,E. Oset and J.R. Pelaez, Phys. Rev Lett. 80,3452(1998).
- [14] T.E.O.Ericson and W. Weise, *Pions and Nuclei*, Clarendon, Oxford (1988).
- [15] C.B.Dover, T.Gutsche, M.Maruyama and A.Faessler, Prog. Part. Nucl. Phys. 29, 87
(1992).
- [16] A.D. Martin and T.D. Spearman,*Elementary Particle Theory*,North-Holland Pub-

- lishing Co.,Amsterdam(1970).
- [17] B.S.Zou and D.V.Bugg, Phys. Rev. D50, 591 (1994);
L.Li, B.S.Zou, G.L.Li, Phys. Rev. D67, 034025 (2003).
 - [18] M.P.Locher, V.E.Markushin, and H.Q.Zheng, Phys. Rev. D55, 2894 (1997).
 - [19] V.Burkert, nucl-th/0005033, in the *Proc. of the NSTAR2000 Conference*, p.1, World Scitific (2001);
Y.Oh, A.I.Titov, and T.S.H.Lee, Phys. Rev. C63, 025201 (2001);
B.Friman and M.Soyeur, Nucl. Phys. A600, 477 (1996).
 - [20] L. Köpke and N.Wermes, Phys Rep. 174, 67 (1989).
 - [21] D.V.Bugg, L.Y.Dong and B.S.Zou, Phys. Lett. B458, 511 (1999).
 - [22] BES Collaboration, Phys. Lett. B440 (1998) 217.
 - [23] BES Collaboration, Phys. Lett. B446 (1999) 356.
 - [24] BES Collaboration, Phys. Lett. B472 (2000) 200.
 - [25] BES Collaboration, Phys. Lett. B472 (2000) 208.
 - [26] BES Collaboration, Phys. Lett. B476 (2000) 25.
 - [27] C.A.Baker et al., Nucl. Phys. A483, 631 (1988);
C. van Eijk et al., Nucl. Phy. A486, 604 (1988);
R. Bacher et al., Z. Phys. A334, 93 (1989).